

FINAL ANSWER KEY

Question 61/2023/OL

Paper Code:

Category 054/2020

Code:

Exam: Assistant Professor Obstetrics and Gynaecology

Date of Test 04-06-2023

Department Government Homoeopathic Medical Colleges

Question1:-The smallest diameter of true pelvis is

A:-Diagonal conjugate

B:-Interspinous diameter

C:-True conjugate

D:-Oblique diameter

Correct Answer:- Option-B

Question2:-The uterine blood flow at term is

A:-50 - 200 ml/min

B:-250 - 450 ml/min

C:-500 - 700 ml/min

D:-800 - 1000 ml/min

Correct Answer:- Option-C

Question3:-Cardiac output during pregnancy is maximal at

A:-36 weeks

B:-32 weeks

C:-28 weeks

D:-24 weeks

Correct Answer:- **Question Cancelled**

Question4:-Pain of uterine contraction is distributed along the following dermatomes

A:-T10 - L1

B:-L2 - L3

C:-L4 - L5

D:-S1 - S2

Correct Answer:- Option-A

Question5:-Commonest type of vertex presentation in an unengaged head is

A:-Left occipito anterior

B:-Right occipito anterior

C:-Left occipito transverse

D:-Right occipito transverse

Correct Answer:- Option-C

Question6:-Commonest diameter of engagement

- I. Mentovertebral diameter
- II. Biparietal diameter
- III. Suboccipitobregmatic
- IV. Suboccipito frontal

A:-Both II and III

B:-Both II and IV

C:-II alone

D:-III alone

Correct Answer:- Option-A

Question7:-About Litzmann Asynclitism

- I. Also called posterior asynclitism
- II. Also called anterior asynclitism
- III. Common in multipara
- IV. Common in primipara

A:-I and III

B:-I and IV

C:-II and III

D:-II and IV

Correct Answer:- Option-B

Question8:-Which is not a surgery for inversion of uterus ?

A:-Haultain

B:-Spincelli

C:-Huntington

D:-Fenton

Correct Answer:- Option-D

Question9:-Death in acute inversion of uterus is due to

A:-Hemorrhage

B:-Neurogenic shock

C:-Infection

D:-Embolism

Correct Answer:- Option-A

Question10:-Role of amniotic fluid turnover is

A:-200 ml/day

B:-300 ml/day

C:-400 ml/day

D:-500 ml/day

Correct Answer:- Option-D

Question11:-pH of amniotic fluid is

A:-6.8

B:-7.0

C:-7.2

D:-7.4

Correct Answer:- Option-C

Question12:-Therapeutic level of magnesium to prevent seizure in eclampsia

A:-1-3 meq/L

B:-4-7 meq/L

C:-8-12 meq/L

D:->12 meq/L

Correct Answer:- Option-B

Question13:-Contraindication for the use of $MgSO_4$ regime in eclampsia

A:-Myasthenia gravis

B:-Bronchial asthma

C:-Tight Mitral Stenosis

D:-Placenta previa

Correct Answer:- Option-A

Question14:-The fetal thyroid begins to produce insulin from

A:-8 weeks

B:-12 weeks

C:-16 weeks

D:-20 weeks

Correct Answer:- Option-B

Question15:-Division of fertilized ovum at 4-8 days result in which type of placentation

A:-DCDA

B:-MCDA

C:-DCMA

D:-MCMA

Correct Answer:- Option-B

Question16:-According to Quintro staging for twin to twin transfusion syndrome, development of hydrops in one twin refers to which stage

A:-Stage II

B:-Stage III

C:-Stage IV

D:-Stage V

Correct Answer:- Option-C

Question17:-Preterm labour is associated mainly with genital tract infection by

A:-Bacterial vaginosis

B:-Trichomonas vaginitis

C:-Candidiasis

D:-Mixed infection

Correct Answer:- Option-A

Question18:-Which type of antigen does not produce erythroblastosis fetalis ?

A:-Duffy

B:-Kell

C:-Kidd

D:-Lewis

Correct Answer:- Option-D

Question19:-Full dose of anti D should be given in a Rh-Negative mother with Rh positive except in

A:-Intra uterine death

B:-After ECV

C:-Spontaneous abortion at 10 weeks

D:-28 weeks of pregnancy

Correct Answer:- Option-C

Question20:-Buddha position of fetus is seen in

A:-Sickle cell anemia

B:-Immune hydrops

C:-Thalassemia

D:-Dimorphic anemia

Correct Answer:- Option-B

Question21:-300 µg of Anti D neutralises _____ fetal blood.

A:-5 mL

B:-10 mL

C:-15 mL

D:-20 mL

Correct Answer:- Option-C

Question22:-First sign of fetal demise is

A:-Robert sign

B:-Halo sign

C:-Spalding sign

D:-Ball sign

Correct Answer:- Option-A

Question23:-Internal podalic version for transverse lie can result in

A:-Cervical laceration

B:-Vaginal laceration

C:-Uterine rupture

D:-Uterine atony

Correct Answer:- Option-C

Question24:-Minimal cervical dialation required for ventouse application

A:-4

B:-6

C:-8

D:-10

Correct Answer:- Option-D

Question25:-Deep transverse arrest is associated with which type of pelvis

A:-Gynecoid

B:-Android

C:-Anthropoid

D:-Pletypelloid

Correct Answer:- Option-B

Question26:-In Mento posteriorpresentation, internal rotation of head occur through

A:-1 /8th circle

B:- 2 /8th circle

C:-3 /8th circle

D:-4 /8th circle

Correct Answer:- Option-C

Question27:-Funnel shaped pelvis is seen in

A:-Tuberculosis

B:-Congenital

C:-Scoliosis

D:-Malignancy

Correct Answer:- Option-A

Question28:-Obstetric conjugate is said to be critical if it is measuring below

A:-8.5 cm

B:-9.0 cm

C:-9.5 cm

D:-10 cm

Correct Answer:- Option-D

Question29:-Vasaprevia is seen in which type of placenta

A:-Central

B:-Velamentous

C:-Battledore

D:-Marginal

Correct Answer:- Option-B

Question30:-Obstructed labour occurs at

A:-Plane of least pelvic dimension

B:-Plane of greatest pelvic dimension

C:-Plane of inlet

D:-Plane of outlet

Correct Answer:- Option-A

Question31:-First sign of uterine rupture

A:-Fetal heart variation

B:-Fresh vaginal blood

C:-Loss of uterine contour

D:-Maternal tachycardia

Correct Answer:- Option-A

Question32:-Worst prognosis of heart disease is seen in

A:-Tight mitral stenosis

B:-Aortic stenosis

C:-Pulmonary artery hypertension

D:-Prosthetic valve

Correct Answer:- Option-C

Question33:-According to Mississippi class system a platelet count of less than 1.5 lakh is classified as

A:-I

B:-II

C:-III

D:-IV

Correct Answer:- Option-C

Question34:-The antidote for magnesium sulfate in eclampsia is

A:-1% calcium gluconate

B:-10% calcium gluconate

C:-1% calcium sulphate

D:-10% calcium sulphate

Correct Answer:- Option-B

Question35:-Vaccine contraindicated in pregnancy

A:-Oral polio

B:-Covid

C:-Influenza

D:-Yellow fever

Correct Answer:- Option-A

Question36:-The risk of development of anencephaly after one affected child is

A:-1%

B:-5%

C:-10%

D:-13%

Correct Answer:- Option-B

Question37:-Nerve roots blocked in pudendal nerve block is

A:-L1, 2, 3

B:-L2, 3

C:-S2, 3, 4

D:-S3, 4

Correct Answer:- Option-C

Question38:-Half life of oxytocin

A:-1-2 min

B:-3-4 min

C:-5-6 min

D:-7-8 min

Correct Answer:- Option-B

Question39:-The first test to become abnormal in iron deficiency anemia

A:-Serum ferritin

B:-Reticulocyte count

C:-Total iron binding capacity

D:-Hematocrit

Correct Answer:- Option-A

Question40:-Following 1 pint blood transfusion, the hematocrit rises by

A:-1%

B:-2%

C:-3%

D:-4%

Correct Answer:- Option-C

Question41:-Single umbilical artery is associated with all except

A:-Abortion

B:-Post maturity

C:-IUGR

D:-Renal anomalies

Correct Answer:- Option-B

Question42:-Transmission of Rubella infection is maximum in

A:-*Ist* trimester

B:-*IInd* trimester

C:-*IIIrd* trimester

D:-In all trimesters

Correct Answer:- Option-A

Question43:-The commonest intrauterine infection in pregnancy is

A:-Cytomegalovirus

B:-Rubella

C:-Toxoplasma

D:-Herpes

Correct Answer:- Option-A

Question44:-Which of the following drug do not cross placenta ?

A:-Warfarin

B:-Heparin

C:-Thyroxine

D:-Methimazole

Correct Answer:- Option-B

Question45:-Reduced maternal serum alpha fetoprotein is seen in

A:-Trisomy 21

B:-IUD

C:-Multiple pregnancy

D:-Neural tube defect

Correct Answer:- Option-A

Question46:-Sinusoidal pattern in cardiotocography denotes

A:-Head compression

B:-Fetal anemia

C:-Placental insufficiency

D:-Loops of cord

Correct Answer:- Option-B

Question47:-Lanugo hair appears at

A:-4 months

B:-5 months

C:-6 months

D:-7 months

Correct Answer:- Option-A

Question48:-Implantation bleeding is also called

A:-Arias-Stella sign

B:-Hartmann sign

C:-Blog sign

D:-Beagle sign

Correct Answer:- Option-B

Question49:-In active phase of labour, the rate of cervical dialation in a primi gravid is

A:-1 cm/hr

B:-1.5 cm/hr

C:-2 cm/hr

D:-2.5 cm/hr

Correct Answer:- Option-A

Question50:-1 pint cryoprecipitate measures

A:-50 ml

B:-100 ml

C:-150 ml

D:-200 ml

Correct Answer:- Option-A

Question51:-True about Ovary is

A:-Attached to uterus by infundibulopelvic ligament

B:-Main blood supply is from Aorta

C:-Does not receive blood supply from uterine artery

D:-Lymphatic drainage is commonly to inguinal lymphnodes

Correct Answer:- Option-B

Question52:-In bimanual palpation

A:-One or two fingers can be inserted into vagina

B:-Normal ovaries are palpable always in lateral fornix

C:-The fundus of the uterus cannot be palpated in a normal uterus

D:-Cervical motion tenderness is normal

Correct Answer:- Option-A

Question53:-Which gynecological cancer has got highest association of hereditary cancer ?

A:-Endometrial Cancer

B:-Ovarian Cancer

C:-Fallopian tube Cancer

D:-Carcinoma Vulva

Correct Answer:- Option-B

Question54:-What is responsible for dramatic increase in local rise in prosta glandins and proteolytic enzymes in follicular wall for ovulation ?

A:-Mid cycle increase in estrogen levels

B:-Midcycle LH surge

C:-The fall in FSH level

D:-Rise in levels of Inhibin B

Correct Answer:- Option-B

Question55:-Most common cause of secondary dysmenorrhoea is

A:-Endometriosis

B:-Fibroids

C:-Nulliparity

D:-PID

Correct Answer:- Option-A

Question56:-In an adoloscent, obese girl with infrequent menstrual cycles and heavy flow, BMI. 25, which investigation is not investigation as first step ?

A:-Serum TSH

B:-Serum Prolactin

C:-Serum Estradiol

D:-Hemoglobin

Correct Answer:- Option-C

Question57:-What is the most effective treatment for vasomotor symptoms in a menopausal lady ?

A:-Isoflavone

B:-Raloxifene

C:-Gabapentin

D:-Estrogen Therapy

Correct Answer:- Option-D

Question58:-Which is wrong regarding Enhanced Recovery After Surgery (ERAS) protocol ?

A:-Patient is allowed to eat ad lib till 6 hours prior to elective surgeries

B:-Patients can take clear fluids by mouth till 2 hours prior to surgery

C:-Patients undergoing pelvic surgery should be kept nil per orally after midnight

D:-More than one class of anti emetic drugs should be used to manage perioperative nausea and vomiting

Correct Answer:- Option-C

Question59:-Schuchardt Incision

A:-is used in myomectomy

B:-used for salpingectomy

C:-may be used in vaginal hysterectomy

D:-may be used in abdominal sacropexy

Correct Answer:- Option-C

Question60:-The recurrence rate of ectopic pregnancy after one ectopic pregnancy

A:-51%

B:-31%

C:-10-15%

D:-3%

Correct Answer:- Option-C

Question61:-Which is a cause of hypergonadotrophic hypogonadism ?

A:-17 α -hydroxylase deficiency in XX individual

B:-Kallman Syndrome

C:-Physiological delay

D:-Central nervous system tumors

Correct Answer:- Option-A

Question62:-Which is not a sign put forward by International Federation for Cervical Pathology and Colposcopy in CIN III ?

A:-Ridge sign

B:-Inner border sign

C:-Rag sign

D:-Rope sign

Correct Answer:- Option-D

Question63:-In a 35 years old P1L1 treatment of choice for CIN 3 is

A:-Colposcopy

B:-LEEP

C:-Cone Biopsy

D:-Watchful expectancy

Correct Answer:- Option-B

Question64:-In Paget's disease of Vulva, which malignancy should be ruled out ?

A:-Adenocarcinoma of anorectum

B:-Squamous carcinoma of Cervix

C:-Endometrioid carcinoma

D:-Immature teratoma of ovary

Correct Answer:- Option-A

Question65:-Which is true about HPV vaccination ?

A:-Maximum protection occurs when girls are given HPV vaccine at 9-15 yrs. of age

B:-HPV vaccination need not be given after 26 years of age

C:-HPV vaccine is only for girls and not for boys

D:-Once a woman is vaccinated with HPV vaccine, there is no role for papsmear or HPV DNA testing

Correct Answer:- Option-A

Question66:-Which is true about "Fertile Window" in a couple trying for conception ?

A:-Fertile window is from Day 10 to Day 21 of the cycle in all women

B:-Fertile window is the 6-day interval ending on the day of ovulation and does not extend post Ovulation

C:-Fertile window is based on concept that Sperms can survive up to only 3 days in well estrogenized cervical mucus

D:-The ovum can survive upto 36 - 48 hours after ovulation

Correct Answer:- Option-B

Question67:-Which is not true regarding POP-Q classification ?

A:-Aa is a point 3cm proximal to external urethral meatus

B:-Bp is in point of maximum prolapse on posterior vaginal wall

C:-C denotes the cervix or cuff

D:-In Procidentia Aa can be - 5cm

Correct Answer:- Option-D

Question68:-Surgical Intervention may not be necessary to evaluate

A:-Ovarian mass greater than 6cm diameter

B:-Adnexal mass greater than 10cm diameter

C:-Solid adnexal mass developing after menopause

D:-Unilocular clear ovarian cyst of 3cm in a perimenopausal lady

Correct Answer:- Option-D

Question69:-Which of the following HPV vaccines is indigenously developed in India ?

A:-CERVAVAC

B:-CERVARIX

C:-Quadrivalent GARDASIL

D:-CERVASIL

Correct Answer:- Option-A

Question70:-Most common degeneration in fibroid is

A:-Red degeneration

B:-Hyaline degeneration

C:-Calcific degeneration

D:-Sarcomatous

Correct Answer:- Option-B

Question71:-Which is true regarding Semen Analysis Criteria (WHO 2010) ?

- i. Volume should be ≥ 1.5 mL
- ii. Sperm motility $> 50\%$ progressive
- iii. White blood cells < 1 million/mL
- iv. Morphology $> 15\%$ normal forms

A:-i and iii are correct

B:-ii and iii are correct

C:-ii and iv are correct

D:-i and iv are correct

Correct Answer:- Option-A

Question72:-Regarding use of Gonadotropins for treatment of infertility in women

- i. In PCOS patients, gonadotropins is associated with increased incidence of OHSS
- ii. Gonadotropins are contraindicated for ovulation induction in women with primary ovarian failure with high FSH
- iii. In certain hypothalamic causes like Kallman's Syndrome, gonadotropins can be used
- iv. Sequential use of gonadotropins and aromatase inhibitors is associated with lower number of follicles

A:-Only i and ii are correct

B:-Only i and iii are correct

C:-i, ii and iii are correct

D:-All are correct

Correct Answer:- Option-D

Question73:-All of the following are indications for myomectomy except

A:-3cm submucous fibroid with recurrent pregnancy loss

B:-5cm submucous fibroid in infertility

C:-6cm subserous fibroid with primary infertility

D:-A lady with 8 cm intramural fibroid in lower segment getting ready for IVF

Correct Answer:- Option-C

Question74:-All of the following are common presentation of adenomyosis except

A:-Secondary infertility

B:-Dysmenorrhoea

C:-Heavy menstrual bleeding

D:-Lower abdominal lump up to umbilicus

Correct Answer:- Option-D

Question75:-All are true regarding gynecological examination of a Pediatric patient except

A:-Young girl can be examined in frog legged position

B:-Young girl can be examined with mother lying on table with legs apart and child on lap with her legs straddling mother's legs

C:-Speculum examination may be performed in pre pubertal girl in office

D:-An Otoscope can be used to examine distal vagina if indicated

Correct Answer:- Option-C

Question76:-All are medical conditions associated with increased risk of osteoporosis after menopause except

A:-Hyperparathyroidism

B:-Hyperthyroidism

C:-Systemic steroid usage

D:-Hormone replacement therapy

Correct Answer:- Option-D

Question77:-The access techniques of laparoscopy in Gynecology are all except

A:-Hasson entry

B:-Pre in sufflation entry

C:-Lateral entry

D:-Direct entry

Correct Answer:- Option-C

Question78:-Operative Hysteroscopy alone may be used often in all except

A:-Removal of a small Figo type 0 myoma

B:-Removal of Figo type 3 myoma

C:-Uterine Septum resection with recurrent pregnancy loss

D:-Removal of missing IUD

Correct Answer:- Option-B

Question79:-All of the following are true except

A:-At birth, cervix and corpus are about equal size

B:-In a reproductive age woman, corpus has grown to two to three times the size of the cervix

C:-"Flexion" is the angle between uterine corpus and upper vagina

D:-After menopause, the transformation zone moves inwards into endocervix

Correct Answer:- Option-C

Question80:-Regarding chemotherapy for malignant ovarian tumors, all are true except

A:-POMB - ACE Regime is for Germ Cell tumors

B:-BEP regimen is for serous tumors

C:-Carboplatin-Paclitaxel combination is for epithelial ovarian cancer

D:-Bevacizumab can be added to platinum drugs in recurrent epithelial ovarian tumors

Correct Answer:- Option-B

Question81:-In a PCOS woman, all are benefits of exercise except

A:-Reduces bodyweight

B:-Reduces insulin resistance

C:-Increases SHBG Levels

D:-Does not have an effect on hirsutism

Correct Answer:- Option-D

Question82:-All are sites of ureteral injury during abdominal hysterectomy except

A:-At pelvic from where ureters cross common iliac artery

B:-In midpelvis as uterine artery crosses over ureter

C:-Near the cervix at the level of internal os

D:-Medial to infundibulopelvic ligament above pelvic brim

Correct Answer:- Option-D

Question83:-Robotic Surgery all are true except

A:-Robotic column holds the robotic arms

B:-For pelvic surgery, the robotic column is side docked over patient's hip

C:-The surgeon sits on patient's right side

D:-Tips of robotic instruments are articulated with seven degrees of freedom

Correct Answer:- Option-C

Question84:-Uterine Artery Embolization (UAE) may be done in all except

A:-8 cm fibroid with pressure symptoms in a 45 years old lady with high risks for anesthesia

B:-7 cm intramural fibroid causing congestive dysmenorrhoea in a 40 yrs old parous lady

C:-32 years old nulliparous lady with 10 cm submucous fibroid

D:-8 cm and 5 cm intramural fibroids on anterior and posterior wall in a 38 years old sterilized women

Correct Answer:- Option-C

Question85:-With metastatic Gestational Trophoblastic Neoplasia, 80% may have visible lung involvement in Chest radiography. All are principal pulmonary patterns except

A:-Alveolar or "snow-storm" pattern

B:-Characteristic B-lines

C:-Pleural effusion

D:-Discrete rounded densities

Correct Answer:- Option-B

Question86:-Treatment of Hirsutism includes all except

A:-Oral contraceptives (COC)

B:-Spironolactone

C:-Norethisterone acetate

D:-Eflornithine HCl

Correct Answer:- Option-C

Question87:-In HIV positive women, all are true except

A:-All HIV infected individuals are recommended to take ART regard less of CD4 T lymphocyte Count

B:-ART is recommended to HIV infected individuals to prevent transmission of infection

C:-Patients with CD4 + T cells count more than 200, should receive prophylaxis against opportunistic infections

D:-CIN progresses to Ca Cervix more rapidly in HIV positive women

Correct Answer:- Option-C

Question88:-Contraindications to medical abortion with MIFEPRISTONE/MISO PROSTOL include all except

A:-Intrauterine device in place

B:-Chronic adrenal failure

C:-Inherited porphyria in woman

D:-Previous history of unsuccessful medical abortion

Correct Answer:- Option-D

Question89:-Regarding Pelvic Inflammatory Disease (PID) all are true except

A:-Out patient treatment is permitted in mild to moderate cases

B:-In diagnosing criteria for PID symptoms need not be present

C:-Women with TO abscess should be treated at hospital

D:-Percutaneous drainage of TO abscess under USG guidance should be tried before starting antibiotics

Correct Answer:- Option-D

Question90:-Regarding laparoscopic Sterilization technique

i. Silastic rubber band, electrocautery, spring clip etc. can be used to occlude fallopian tube.

ii. Reversal of sterilization is more successful following falope ring than electrocoagulation.

iii. Unipolar electrocoagulation is recommended as it is associated with lesser complications.

A:-i and ii are true

B:-i, ii and iii are true

C:-i and iii are true

D:-i alone is true

Correct Answer:- Option-A

Question91:-In Genital Tuberculosis which is the most common part affected in a woman ?

A:-Endometrium

B:-Fallopian tube

C:-Ovary

D:-Pelvic peritoneum

Correct Answer:- Option-B

Question92:-Which is contraindication for medical management of ectopic pregnancy with Methotrexate ?

i. Breast feeding

ii. Severe anemia

iii. 5 cm diameter ectopic mass by TVS

iv. β HCG a value of 4000 mIU/mL

A:-i and ii

B:-i, ii and iii

C:-i and iv

D:-ii and iv

Correct Answer:- Option-B

Question93:-In surgical prevention of hereditary cancers in gynecology

i. Gold standard and most effective method of preventing ovarian cancer in high risk women is Risk Reducing Salpingo Oophorectomy (RRSO)

ii. RRSO reduces ovarian cancer risk by 80-96% in BRCA1/BRCA2 Carriers

iii. RRSO is useful in BRCA1/BRCA2 and Lynch syndrome in almost similar degrees

A:-i and ii are true

B:-i and iii are true

C:-All are true

D:-Only ii is true

Correct Answer:- Option-A

Question94:-About Condoms which is true

A:-Risk of condom breakage is 13% and mainly due to friction

B:-K-Y jelly can be used to increase the lubrication

C:-Condoms do not reduce the risk of invasive Cervical Cancer

D:-Polyurethane and Tactylon condoms should be offered to men with latex allergy

Correct Answer:- Option-D

Question95:-In different methods to extract sperm from Seminal plasma, in IUI which method yields the best motile sperms ?

A:-Washing

B:-Swim up

C:-Density gradient centrifugation

D:-Serial filtration

Correct Answer:- Option-C

Question96:-Regarding imaging in Gynecology which is not true

A:-Trans vaginal ultrasound transducer operates at 5 – 8 MHz

B:-Trans abdominal ultrasound transducer operates at 2.5 – 3.5 MHz and hence has got better resolution for normal sized ovaries within pelvis

C:-MRI is better than CT Scan in evaluating soft Tissue involvement in Ca Endometrium

D:-Saline infusion sonography helps to differentiate sessile sub mucus fibroid and endometrial polyp

Correct Answer:- Option-B

Question97:-Regarding Vulvovaginal Candidiasis (VVC) which is not true ?

A:-80% of women would have at least one episode in life time

B:-Recurrent VVC is defined as more than 4 episodes in a year

C:-Treatment of recurrent VVC include oral Fluconazole 150 mgs every three days for 3 doses and suppressive 150 mg weekly for 6 months

D:-pH of vagina in women with recurrent VVC is 6 or above

Correct Answer:- Option-D

Question98:-Which is not true regarding classification of Female Genital Mutation (FGM) ?

A:-Type I - Removing all or part of clitoris and prepuce

B:-Type III - Narrowing vaginal orifice and creating a covering of adjacent Labia Majora and Minora

C:-Type II - Excision of part or all of clitoris and labia minora with or without excision of Labia majora

D:-Type IV - Radical procedure involving removal of clitoris, Labia majora and Labia minora leaving an opening

Correct Answer:- Option-D

Question99:-Regarding tubal recanalization which is not true

A:-main predictors of success are age < 35 ys, isthmic - isthmic, anastomosis

B:-IVF is better option when there are severe pelvic adhesions

C:-Pregnancy rates following microsurgical tubal reanastomosis for sterilization reversal is 55 - 81%

D:-Success rate is almost similar when done for Hydrosalpinx correction

Correct Answer:- Option-D

Question100:-Which is not a Surgical procedure described for Stress Urinary Incontinance (SUI) ?

A:-Bladder neck suspension

B:-Mid Urethral Sling

C:-Distal Urethral Plication Stitch

D:-Kelly's Stitch

Correct Answer:- Option-C